



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION  
International Bureau

## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup>:</b> A61K 38/00, 39/00, C07H 21/04, C12N 1/19, 1/21, 5/10, 15/00, 15/70, 15/79, C12P 21/06	<b>A1</b>	<b>(11) International Publication Number:</b> WO 95/04542 <b>(43) International Publication Date:</b> 16 February 1995 (16.02.95)
<b>(21) International Application Number:</b> PCT/US94/08721 <b>(22) International Filing Date:</b> 2 August 1994 (02.08.94)  <b>(30) Priority Data:</b> 08/103,623 6 August 1993 (06.08.93) US  <b>(60) Parent Application or Grant</b> <b>(63) Related by Continuation</b> US 08/103,623 (CON) Filed on 6 August 1993 (06.08.93)  <b>(71) Applicant (for all designated States except US):</b> CYTEL CORPORATION [US/US]; 3525 John Hopkins Court, San Diego, CA 92121 (US).  <b>(72) Inventors; and</b> <b>(75) Inventors/Applicants (for US only):</b> FIKES, John, D. [US/US]; 6474 Lipmann Street, San Diego, CA 92122 (US). LIV-INGSTON, Brian, D. [US/US]; 8615 Covina Street, San Diego, CA 92126 (US). SETTE, Alessandro, D. [IT/US]; 5551 Linda Rosa Avenue, La Jolla, CA 92037 (US). SIDNEY, John, C. [US/US]; 8541D Villa La Jolla Drive, La Jolla, CA 92037 (US).		<b>(74) Agents:</b> PARMELEE, Steven, W. et al.; Townsend and Townsend Khourie and Crew, One Market Plaza, 20th floor, Steuart Street Tower, San Francisco, CA 94105 (US).  <b>(81) Designated States:</b> AM, AT, AU, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, ES, FI, GB, GE, HU, JP, KE, KG, KP, KR, KZ, LK, LT, LU, LV, MD, MG, MN, MW, NL, NO, NZ, PL, PT, RO, RU, SD, SE, SI, SK, TJ, TT, UA, US, UZ, VN, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG), ARIPO patent (KE, MW, SD).  <b>Published</b> <i>With international search report.</i>
<b>(54) Title:</b> CLONING AND CHARACTERIZATION OF THE COMPLETE MAGE-1 GENE		
<b>(57) Abstract</b>  The complete nucleotide and amino acid sequences of the human MAGE-1 antigen are provided. Peptides from residues of the C-terminal are used to define epitopes that stimulate HLA-restricted cytotoxic T lymphocyte activity against MAGE-1 antigens. The peptides are particularly useful in methods for stimulating the immune response of individuals against MAGE-1 antigens associated with melanomas.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

## 5 CLONING AND CHARACTERIZATION OF THE COMPLETE MAGE 1 GENE

10 Background of the Invention

Tumor cells express certain antigens which are absent from, or present in small amounts on, their normal cellular counterparts. Most of these are differentiation antigens, shared by the tumor and various embryonic cells. 15 Some of the antigens that appear with sufficient selectivity may serve as possible targets for therapeutic agents.

More than 40 different melanoma antigens have been defined with monoclonal antibodies, resulting in several major antigenic families with immunologically and biologically 20 distinct characteristics. Among these families are: (1) the high molecular weight oncofetal proteins; (2) the gangliosides; (3) receptors for growth factors such as EGF, PDGF, TGF-alpha and TGF-beta, and nerve growth factor; (4) cation transport and binding proteins such as p97; (5) HLA 25 class II antigens; (6) pigmentation-associated antigens; and (7) extracellular matrix proteins. Herlyn and Koproski, Ann. Rev. Immunol. 6: 283-308 (1988).

While preliminary studies with monoclonal antibody-based therapy and diagnosis directed toward various of these 30 antigens are encouraging, work continues unabated in the hope that better agents and antigenic targets can be identified. Cutaneous malignant melanoma is increasing in prevalence at an alarming rate, particularly in the United States.

More recently a new family of antigens has been 35 described on melanoma tumors. These antigens, now termed the "melanoma antigen," or MAGE family of antigens, were identified in a melanoma cell line which was lysed by a panel of autologous cytotoxic T lymphocytes ("CTLs"). Cells which did not express a MAGE-type antigen were not killed by the 40 CTL, and by selecting these "antigen-loss" variants, six independent antigens were identified. Van den Eynde et al., Int. J. Cancer, 44: 634 (1989). A gene encoding one of the antigens, designated MZ2-E ("E"), has been cloned and

sequenced. Van der Bruggen et al., Science 254: 1643 (1991). The sequence was deposited in GenBank (accession #M77481), and comparison of the nucleotide sequence, designated "MAGE-1," failed to reveal any significant homology with any sequence in data banks, including GenBank. Two additional nonidentical cDNAs were also found (MAGE-2 and MAGE-3) which were more closely related to each other than to MAGE-1, but the three were approximately equally expressed.

Smaller regions of the MAGE-1 gene were cloned and transfected into cells. These transfectants expressed antigen which was recognized by the anti-E CTLs. Thus, it appears that the gene does not encode a protein which further activates an antigen-encoding gene. Van der Bruggen, id. The sequence encoding the antigenic peptide was speculated to be within the region of overlap of the segments. See Traversari et al., J. Exp. Med. 176: 1453-1457 (1992). The cDNAs of MAGE-2 and MAGE-3 were unable to transfer the expression of antigen E in transfection experiments. The presenting molecule for the E-antigen was thought to be HLA-A1.

The MAGE gene family has been shown by Van der Bruggen et al., id., to be expressed by a variety of different tumors and are not limited to melanomas, but they are not expressed by most normal cells. Thus, the MAGE antigens may have important implications for cancer immunotherapy. The sequence of the MAGE-1 gene was thought to be identical in both normal tissues and in tumors.

What is needed in the art is a more thorough understanding of the immunogenic tumor-rejecting epitopes of the MAGE antigens. Once the immunodominant epitopes are identified, along with their HLA restriction, more effective therapeutic protocols can be devised. The present invention fulfills these and other related needs.

#### Summary of the Invention

This invention is based in part on the novel and unexpected observation that the previously reported gene encoding the human MAGE-1 protein encodes an additional fifty-eight amino acids at the C-terminal end. The complete human

MAGE-1 protein and peptides thereof can be produced by recombinant or synthetic means and may or may not have the biological activity of the native MAGE-1 antigen, depending on the intended use. Accordingly, isolated and purified polynucleotides are described which code for the complete human MAGE-1 protein. The cDNA which codes for the full length human MAGE-1 protein may be incorporated into a recombinant DNA vector; which in turn may be used to transform a suitable host; the host cell transformed with the vector including the cDNA can express full length human MAGE-1 protein, and the full length human MAGE-1 protein can be recovered.

This invention further concerns MAGE-1 immunogenic peptides from the C-terminus of the MAGE-1 protein which induce CTL activity. The immunogenic peptides of this invention may be identified using motifs as described in copending U.S. patent applications S.N. 07/926,666 and S.N. 08/027,146 for the various MHC class I alleles. Thus, small synthetic or recombinant peptides can be prepared which immunologically mimic MAGE-1 CTL inducing antigenic determinants. The CTL-inducing MAGE-1 peptides of the invention can be used therapeutically, for example, to induce, in the context of an appropriate MHC presenting molecule, an immunological response to tumors which express the corresponding MAGE determinants. In this manner the tumor cells can be killed or inhibited. The induction of CTLs can be accomplished in vivo or ex vivo. Thus, the MAGE-1 peptides described herein also can be formulated and administered as pharmaceutical compositions, especially when used to induce immunological responses in individuals predisposed to developing or already afflicted by a tumor which expresses MAGE-1 determinants.

In yet other embodiments the invention relates to methods for diagnosis, where the peptides of the invention are used to determine the presence in an individual of lymphocytes which are capable of a cytotoxic T cell response to MAGE-1 antigen. Typically the lymphocytes are peripheral blood lymphocytes and the individual of interest is suffering from a

tumor associated with MAGE antigen. The diagnostic methods and compositions can be used in conjunction with therapeutic approaches to MAGE related diseases, and particularly the treatment of malignant melanoma.

5

#### Brief Description of the Drawings

Fig. 1 is the nucleotide and amino acid sequence of full length human MAGE-1 protein.

10

Fig. 2 is the nucleotide and amino acid sequence of the newly discovered C terminal portion of the full length human MAGE-1 protein.

15

Fig. 3 illustrates CTL responses specific for certain newly identified peptides from the C-terminus portion of the human MAGE-1 protein.

#### Description of the Specific Embodiments

20

The melanoma antigen termed "MAGE" was identified in the context of a CTL-inducing antigen. The MAGE antigen has since been discovered to be a family of related antigens expressed by a variety of tumor cells. The present invention provides the complete nucleotide sequence encoding human MAGE-1 antigen and the complete amino acid sequence thereof, thereby providing for the ultimate expression of the complete human MAGE-1 protein and new MAGE-1 peptides which have immunological activity. Recombinant DNA expression systems and chemical synthetic methods provide convenient means for obtaining large quantities of recombinant human MAGE-1 and the peptide fragments thereof in relatively pure form.

25

30

In preferred embodiments the peptides of the invention are derived from the region of the MAGE-1 antigen of the C-terminal 58 amino acids, as set forth in Seq. ID No. 1:

35

Seq. ID No. 1

Arg-Gln-Val-Pro-Asp-Ser-Asp-Pro-Ala-Arg-Tyr-Glu-Phe-Leu-Trp-Gly-Pro-Arg-Ala-Leu-Ala-Glu-Thr-Ser-Tyr-Val-Lys-Val-Leu-

Glu-Tyr-Val-Ile-Lys-Val-Ser-Ala-Arg-Val-Arg-Phe-Phe-Phe-Pro-Ser-Leu-Arg-Glu-Ala-Ala-Leu-Arg-Glu-Glu-Glu-Glu-Gly-Val

5       The peptide which is selected from the region of  
Seq. ID No. 1 induces MHC HLA class I-restricted CTL responses  
to MAGE expressing cells. The stimulated CTL, which secrete  
lymphokines (e.g., gamma interferon) and liberate products  
(e.g., proteolytic enzymes such as serine esterases) that  
10       inhibit viral replication in infected autologous cells or  
transfected cells, with or without cell killing, are able to  
interrupt or substantially prevent the growth of MAGE  
expressing tumor cells. In many instances the combination of  
an effective cytotoxic T cell response and a protective  
antibody response to selected tumor antigens will be preferred  
15       for treating a MAGE-associated tumor.

      In more preferred embodiments described herein an  
immunity-inducing peptide derived from the region of Seq. ID  
No. 1 has at least seven amino acids wherein a majority of  
amino acids of the peptide will be identical or substantially  
20       homologous, when compared to the amino acids comprising the  
corresponding portion of the naturally occurring MAGE-1  
sequence. Representative peptides of this region are set  
forth in Table 1 below with MHC restriction indicated.

      The peptide can be optionally flanked and/or  
25       modified at one or both of the N- and C-termini, as desired,  
by amino acids from MAGE sequences, particularly MAGE-1, amino  
acids added to facilitate linking, other N- and C-terminal  
modifications, linked to carriers, etc., as further described  
herein. The peptide induces a CTL response which is mediated  
30       by at least the MHC class I molecule as indicated above.

      The term "peptide" is used interchangeably with  
"oligopeptide" in the present specification to designate a  
series of residues, typically L-amino acids, connected one to  
the other typically by peptide bonds between the alpha-amino  
35       and carbonyl groups of adjacent amino acids. The  
oligopeptides of the invention are less than about 15 residues  
in length and usually consist of between about 8 and about 11  
residues, preferably 9 or 10 residues.

By "immunogenic peptide" of the present invention is meant a peptide which comprises an allele-specific motif such that the peptide will bind the MHC allele and be capable of inducing a CTL response. The immunogenic peptides of this invention are derived from selected epitopic regions of the C-terminal 58 amino acid residues of the MAGE-1 antigen. The immunogenic peptides are capable of binding to an appropriate class I MHC molecule and inducing a cytotoxic T cell response against the MAGE antigen from which the immunogenic peptide is derived.

A "conserved residue" is an amino acid which occurs in a significantly higher frequency than would be expected by random distribution at a particular position in a peptide motif. Typically a conserved residue is one at which the immunogenic peptide may provide a contact point with the MHC molecule. One to three, preferably two, conserved residues within a peptide of defined length defines a motif for an immunogenic peptide. These residues are typically in close contact with the peptide binding groove, with their side chains buried in specific pockets of the groove itself. Typically, an immunogenic peptide will comprise up to three conserved residues, more usually two conserved residues.

As used herein, "negative binding residues" are amino acids which if present at certain positions will result in a peptide being a nonbinder or poor binder and in turn fail to induce a CTL response despite the presence of the appropriate conserved residues within the peptide.

The term "motif" refers to the pattern of residues in a peptide of defined length, usually about 8 to about 11 amino acids, which is recognized by a particular MHC allele. The peptide motifs are typically different for each human MHC allele and differ in the pattern of the highly conserved residues.

The binding motif for an allele can be defined with increasing degrees of precision. In one case, all of the conserved residues are present in the correct positions in a peptide and there are no negative binding residues present.



The phrases "isolated" or "biologically pure" refer to material which is substantially or essentially free from components which normally accompany it as found in its native state. Thus, the peptides of this invention do not contain materials normally associated with their in situ environment, e.g., MHC I molecules on antigen presenting cells. Even where a protein has been isolated to a homogenous or dominant band, there are trace contaminants in the range of 5-10% of native protein which co-purify with the desired protein. Isolated peptides of this invention do not contain such endogenous co-purified protein.

The term "residue" refers to an amino acid or amino acid mimetic incorporated in a oligopeptide by an amide bond or amide bond mimetic.

Peptides comprising the CTL epitopes are synthesized and the ability to bind appropriate MHC molecules is determined in assays using, for example, purified class I molecules and radioiodinated peptides and/or cells expressing empty class I molecules by, for instance, immunofluorescent staining and flow microfluorimetry, peptide-dependent class I assembly assays, and inhibition of CTL recognition by peptide competition. Those peptides that bind to the class I molecule are further chosen for their ability to serve as targets for CTLs derived from afflicted individuals, as well as for their capacity to induce primary in vitro or in vivo CTL responses that can give rise to CTL populations capable of reacting with tumor cells as therapeutic agents. Methods for determining allele-specific peptides and peptide motifs are described in co-pending commonly owned applications USSN 027,146 and USSN 027,746, which are incorporated herein by reference.

The peptides or oligopeptides can be prepared "synthetically," as described herein below, or by recombinant DNA technology. Although the peptide will preferably be substantially free of other naturally occurring human proteins and fragments thereof, in some embodiments the peptides can be conjugated to other MAGE fragments or other proteins or peptides which contribute directly or indirectly to an anti-tumor immunological response. The term peptide or

oligopeptide is used interchangeably with polypeptide in the present specification to designate a series of amino acids connected one to the other by peptide bonds between the alpha-amino and alpha-carboxy groups of adjacent amino acids.

5 The polypeptides or peptides can be a variety of lengths, either in their neutral (uncharged) forms or in forms which are salts, and either free of modifications such as glycosylation, side chain oxidation, or phosphorylation or containing these modifications, subject to the condition that  
10 the modification not destroy the biological activity of the polypeptides as herein described.

Desirably, the peptide will be as small as possible while still maintaining substantially all of the biological activity of the large peptide. When possible, it may be  
15 desirable to optimize peptides of the invention to a length of nine or ten amino acid residues, commensurate in size with processed peptides that are bound to MHC class I molecules on the cell surface. See generally, Schumacher et al., Nature 350:703-706 (1991); Van Bleek et al., Nature 348:213-216  
20 (1990); Rotzschke et al., Nature 348:252-254 (1990); and Falk et al., Nature 351:290-296 (1991), which are incorporated herein by reference. By biological activity is meant the ability to bind an appropriate MHC molecule and, in the case of peptides useful for stimulating CTL responses, induce a CTL  
25 response against MAGE antigen or antigen mimetic. In the case of a peptide analog antagonist, the analog will have biological activity if it competes with the peptide for binding to the MHC molecule and has a substantially reduced ability to stimulate a CTL response when compared to the  
30 native peptide. By a CTL response is meant a CD8<sup>+</sup> T lymphocyte response specific for the MAGE antigen of interest, e.g., members of the MAGE antigen family, wherein CD8<sup>+</sup>, MHC class I-restricted T lymphocytes are activated. As noted above, the activated T lymphocytes will secrete a variety of  
35 products which inhibit tumor cell replication and may or may not directly kill the tumor cell or other transfected cell which expresses the appropriate MAGE antigenic determinant(s).

The terms "homologous", "substantially homologous", and "substantial homology" as used herein denote a sequence of amino acids having at least 50% identity wherein one sequence is compared to a reference sequence of amino acids. The percentage of sequence identity or homology is calculated by comparing one to another when aligned to corresponding portions of the reference sequence.

The peptides of the present invention or analogs thereof which have CTL stimulating activity may be modified to provide desired attributes other than improved serum half life. For instance, the ability of the peptides to induce CTL activity can be enhanced by linkage to a sequence which contains at least one epitope that is capable of inducing a T helper cell response. Particularly preferred immunogenic peptides/T helper conjugates are linked by a spacer molecule. The spacer is typically comprised of relatively small, neutral molecules, such as amino acids or amino acid mimetics, which are substantially uncharged under physiological conditions. The spacers are typically selected from, e.g., Ala, Gly, or other neutral spacers of nonpolar amino acids or neutral polar amino acids. It will be understood that the optionally present spacer need not be comprised of the same residues and thus may be a hetero- or homo-oligomer. When present, the spacer will usually be at least one or two residues, more usually three to six residues. Alternatively, the CTL peptide may be linked to the T helper peptide without a spacer.

The immunogenic peptide may be linked to the T helper peptide either directly or via a spacer either at the amino or carboxy terminus of the CTL peptide. The amino terminus of either the immunogenic peptide or the T helper peptide may be acylated.

In some embodiments it may be desirable to include in the pharmaceutical compositions of the invention at least one component which assists in priming CTL. Lipids have been identified as agents capable of assisting the priming CTL in vivo against certain antigens. For example, palmitic acid residues can be attached to the alpha and epsilon amino groups of a Lys residue and then linked, e.g., via one or more

linking residues such as Gly, Gly-Gly-, Ser, Ser-Ser, or the like, to an immunogenic peptide. The lipidated peptide can then be injected directly in a micellar form, incorporated into a liposome or emulsified in an adjuvant, e.g., incomplete Freund's adjuvant. In a preferred embodiment a particularly effective immunogen comprises palmitic acid attached to alpha and epsilon amino groups of Lys, which is attached via linkage, e.g., Ser-Ser, to the amino terminus of the immunogenic peptide.

As another example of lipid priming of CTL responses, *E. coli* lipoproteins, such as tripalmitoyl-S-glycerylcysteinylserine (P<sub>3</sub>CSS) can be used to prime specific CTL when covalently attached to an appropriate peptide. See, e.g., Deres et al., *Nature* 342:561-564 (1989), incorporated herein by reference. Peptides of the invention can be coupled to P<sub>3</sub>CSS, for example, and the lipopeptide administered to an individual to specifically prime a CTL response to the target antigen. Further, as the induction of neutralizing antibodies can also be primed with P<sub>3</sub>CSS conjugated to a peptide which displays an appropriate epitope, the two compositions can be combined to more effectively elicit both humoral and cell-mediated responses to MAGE antigen.

As mentioned above, additional amino acids can be added to the termini of an oligopeptide or peptide to provide for ease of linking peptides one to another, for coupling to a carrier, support or larger peptide, for reasons discussed herein; for modifying the physical or chemical properties of the peptide or oligopeptide, or the like. Amino acids such as tyrosine, cysteine, lysine, glutamic or aspartic acid, or the like, can be introduced at the C- or N-terminus of the peptide or oligopeptide. In addition, the peptide or oligopeptide sequences can differ from the natural sequence by being modified by terminal-NH<sub>2</sub> acylation, e.g., by alkanoyl (C<sub>1</sub>-C<sub>20</sub>) or thioglycolyl acetylation, terminal-carboxy amidation, e.g., ammonia, methylamine, etc. In some instances these modifications may provide sites for linking to a support or other molecule.

It will be understood that the peptides of the present invention or analogs thereof which have CTL and/or T helper stimulating activity may be modified to provide other desired attributes, e.g., improved pharmacological characteristics, while increasing or at least retaining substantially all of the biological activity of the unmodified peptide. For instance, the peptides can be modified by extending, decreasing or substituting in the peptides amino acid sequences by, e.g., the addition or deletion of amino acids on either the amino terminal or carboxy terminal end, or both, of peptides derived from the sequences disclosed herein. The CTL activity of the subject peptides can be enhanced by linkage to a sequence which contains at least one epitope that is capable of inducing a T helper cell response, as explained above.

The peptides employed in the subject invention need not be identical to those exemplary peptides identified above or to a particular MAGE or MAGE-1 protein sequence, so long as the subject compounds are able to bind to the appropriate MHC molecule and provide for cytotoxic T lymphocytic or T helper activity against cells which express a MAGE antigen. Therefore, the peptides may be subject to various changes, such as insertions, deletions, and substitutions, either conservative or non-conservative, where such changes might provide for certain advantages in their use. By conservative substitutions is meant replacing an amino acid residue with another which is biologically and/or chemically similar, e.g., one hydrophobic residue for another, or one polar residue for another. The substitutions include combinations such as Gly, Ala; Val, Ile, Leu; Asp, Glu; Asn, Gln; Ser, Thr; Lys, Arg; and Phe, Tyr. Usually, the portion of the sequence which is intended to substantially mimic a MAGE CTL or T helper stimulating epitope will not differ by more than about 20% from the sequence of at least one member of the MAGE family, except where additional amino acids may be added at either terminus for the purpose of modifying the physical or chemical properties of the peptide for, e.g., ease of linking or coupling, and the like. In those situations where regions of

the peptide sequences are found to be polymorphic among MAGE antigens, it may be desirable to vary one or more particular amino acids to more effectively mimic differing cytotoxic T-lymphocyte or T helper epitopes of different MAGE antigens.

5           Using the methods described herein, two or more peptides may be identified which define different or overlapping CTL or T helper epitopes from a particular region. For example, using the methods described herein, two or more peptides may define different or overlapping CTL or T helper  
10           epitopes from a particular region, e.g., the peptide region of the C-terminus of MAGE-1, or of a different region, which peptides can be combined in a "cocktail" to provide enhanced immunogenicity for CTL or T helper-mediated responses. Peptides of one region can also be combined with peptides  
15           having different MHC restriction elements. This composition can be used to effectively broaden the immunological coverage provided by therapeutic, vaccine or diagnostic methods and compositions of the invention among a diverse population. When the peptides are linked, by covalent or non-covalent  
20           means, it will be understood that linkage should not substantially interfere with either of the linked groups to function as described, e.g., to function as a MAGE cytotoxic T cell determinant or MAGE T helper determinant.

          In another aspect the peptides of the invention can  
25           be combined or coupled with other peptides which present MAGE T helper cell epitopes, i.e., T helper peptides comprising six to thirty amino acids containing a T helper epitope from a MAGE protein or other immunogenic protein or derivative thereof to stimulate T cells that cooperate in the induction  
30           of immune response against the MAGE antigen, e.g., in the CTL response to MAGE determinants. The T-helper cells can be either the T-helper 1 or T-helper 2 phenotype, for example. Compositions of T-helper peptides and CTL peptides thereby enhance an individual's immunity by providing cell-mediated  
35           immunity and protective antibodies to MAGE antigen. T helper epitopes can be provided by peptides from, for example, tetanus toxoid<sub>830-843</sub> having the sequence Gln-Tyr-Ile-Lys-Ala-Asn-Ser-Lys-Phe-Ile-Gly-Ile-Thr-Glu (QYIKANSKFIGITE) [Seq. ID

No. 18]; malaria circumsporozoite<sub>382-398</sub> Lys-Ile-Ala-Lys-Met-Lys-Ala-Ser-Ser-Val-Phe-Asn-Val-Val-Asn-Ser (KIAKMEKASSVFNVVNS) [Seq. ID No. 19]; malaria circumsporozoite<sub>378-398</sub> Asp-Ile-Glu-Lys-Lys-Ile-Ala-Lys-Met-Lys-Ala-Ser-Ser-Val-Phe-Asn-Val-Val-Asn-Ser (DIEKKIAKMEKASSVFNVVNS) [Seq. ID No. 20]; ovalbumin<sub>323-336</sub> Ile-Ser-Gln-Ala-Val-His-Ala-Ala-His-Ala-Glu-Ile-Asn-Glu [Seq. ID No. 21], the influenza epitope<sub>307-319</sub> Pro-Lys-Tyr-Val-Lys-Gln-Asn-Thr-Leu-Lys-Leu-Ala-Thr [Seq. ID No. 22], and others.

In preferred embodiments the CTL inducing peptides of the invention are covalently linked to the T helper peptides. Particularly preferred CTL inducing peptides / T helper conjugates are linked by a spacer molecule. Alternatively, the CTL peptide may be linked to the T helper peptide without a spacer. The T helper peptide is conjugated to the CTL peptide, preferably with the T helper peptide positioned at the amino terminus. The peptides may be joined by a neutral linker, such as Ala-Ala-Ala or the like, and preferably further contains a lipid residue such as palmitic acid or the like which is attached to alpha and epsilon amino groups of a Lys residue ((PAM)<sub>2</sub>Lys), which is attached to the amino terminus of the peptide conjugate, typically via Ser-Ser linkage or the like.

The peptides of the invention can be prepared in a wide variety of ways. Because of their relatively short size, the peptides can be synthesized in solution or on a solid support in accordance with conventional techniques. Various automatic synthesizers are commercially available and can be used in accordance with known protocols. See, for example, Stewart and Young, Solid Phase Peptide Synthesis, 2d. ed., Pierce Chemical Co. (1984); Tam et al., J. Am. Chem. Soc. 105:6442 (1983); Merrifield, Science 232:341-347 (1986); and Barany and Merrifield, The Peptides, Gross and Meienhofer, eds., Academic Press, New York, pp. 1-284 (1979), each of which is incorporated herein by reference.

Alternatively, recombinant DNA technology may be employed wherein a nucleotide sequence which encodes a CTL peptide and/or T helper peptide of interest is inserted into

an expression vector, transformed or transfected into an appropriate host cell and cultivated under conditions suitable for expression. These procedures are generally known in the art, as described generally in Sambrook et al., Molecular Cloning, A Laboratory Manual, Cold Spring Harbor Press, Cold Spring Harbor, New York (1982), and Ausubel et al., (ed.) Current Protocols in Molecular Biology, John Wiley and Sons, Inc., New York (1987), and U.S. Pat. Nos. 4,237,224, 4,273,875, 4,431,739, 4,363,877 and 4,428,941, for example, which disclosures are incorporated herein by reference. Thus, fusion proteins which comprise one or more peptide sequences of the invention can be used to present the MAGE-1 CTL determinants. For example, a recombinant MAGE antigen polypeptide is prepared in which the amino acid sequence is altered so as to more effectively present epitopes of peptide regions described herein to stimulate a CTL response.

As the coding sequence for peptides of the length contemplated herein can be synthesized by chemical techniques, for example, the phosphotriester method of Matteucci et al., J. Am. Chem. Soc. 103:3185 (1981), modification can be made simply by substituting the appropriate base(s) for those encoding the native peptide sequence. The coding sequence can then be provided with appropriate linkers and ligated into expression vectors commonly available in the art, and the vectors used to transform suitable hosts to produce the desired fusion protein. A number of such vectors and suitable host systems are now available. For expression, the coding sequence will be provided with operably linked start and stop codons, promoter and terminator regions and usually a replication system to provide an expression vector for expression in the desired cellular host. Of course, bacterial, yeast or mammalian cell hosts may be used, employing suitable vectors and control sequences.

The complete MAGE-1 DNA sequence or fragments encoding the C-terminal 58 amino acids of the MAGE-1 protein as described herein may be introduced into cultured mammalian cells by a variety of means, as will be recognized by those skilled in the art. For example, calcium phosphate-mediated



transfection (Wigler et al., Cell 14: 725, 1978; Corsaro and Pearson, Somatic Cell Genetics 7: 603, 1981; Graham and Van der Eb, Virology 52: 456, 1973), electroporation (Neumann et al., EMBO J. 1: 841-845, 1982), or DEAE-dextran mediated  
5 transfection (Ausubel et al., (ed.) Current Protocols in Molecular Biology, John Wiley and Sons, Inc., NY (1987), incorporated herein by reference) may find convenient use. To identify cells that have stably integrated the cloned DNA, a selectable marker is generally introduced into the cells along  
10 with the gene or cDNA of interest. Preferred selectable markers for use in cultured mammalian cells include genes that confer resistance to drugs, such as neomycin, hygromycin, and methotrexate. Further, the selectable marker may be an amplifiable selectable marker, and preferred amplifiable  
15 selectable markers include the DHFR gene and the neomycin resistance gene. Selectable markers are reviewed by Thilly (Mammalian Cell Technology, Butterworth Publishers, Stoneham, MA, which is incorporated herein by reference).

Transfected mammalian cells are allowed to grow for  
20 a period of time, typically 1-2 days, to begin expressing the MAGE-1 DNA sequence(s) of interest. Drug selection is then applied to select for growth of cells that are expressing the selectable marker in a stable fashion. For cells that have been transfected with an amplifiable selectable marker the  
25 drug concentration may be increased in a stepwise manner to select for increased copy number of the cloned sequences, thereby increasing expression levels.

Promoters, terminators and methods for introducing expression vectors encoding foreign proteins such as human  
30 MAGE-1 into plant, avian and insect cells are also well known in the art. Techniques for transforming fungi are well known in the literature, and have been described, for instance, by Beggs (Nature 275:104-108 (1978)), Hinnen et al. (Proc. Natl. Acad. Sci. USA 75: 1929-1933, 1978), Yelton et al. (Proc.  
35 Natl. Acad. Sci. USA 81: 1740-1747, 1984), Russell (Nature 301: 167-169, 1983) and U.S. Patent 4,935,349, incorporated herein by reference. Suitable yeast vectors for use in the present invention will generally include a selectable marker,

which may be one of any number of genes that exhibit a dominant phenotype for which a phenotypic assay exists to enable transformants to be selected.

Host cells containing DNA constructs which encode the complete MAGE-1 protein of the present invention or C-terminal fragments thereof are then cultured to produce the protein or peptides. The cells are cultured according to standard methods in a culture medium containing nutrients required for growth of the chosen host cells. A variety of suitable media are known in the art. The growth medium will generally select for cells containing the DNA construct by, for example, drug selection or deficiency in an essential nutrient which is complemented by the selectable marker on the DNA construct or co-transfected with the DNA construct. Selection of a medium appropriate for the particular cell line used is within the level of ordinary skill in the art. Cultured mammalian cells are generally cultured in commercially available serum-containing or serum-free media.

The complete MAGE-1 protein and C-terminal fragments thereof produced according to the present invention may be purified by, e.g., affinity chromatography on an antibody column using antibodies, preferably monoclonal antibodies, directed against corresponding MAGE epitopes. Additional purification may be achieved by conventional chemical purification means, such as liquid chromatography, gradient centrifugation, and gel electrophoresis, among others. Methods of protein purification are known in the art (see generally, Scopes, R., Protein Purification, Springer-Verlag, NY (1982), which is incorporated herein by reference) and may be applied to the purification of the recombinant human MAGE-1 described herein. Substantially pure recombinant human MAGE-1 of at least about 50% is preferred, at least about 70-80% more preferred, and 95-99% or more homogeneity most preferred, particularly for pharmaceutical uses. Once purified, partially or to homogeneity, as desired, the recombinant human MAGE may then be used diagnostically, therapeutically, etc. as described herein.

The peptides of the present invention and pharmaceutical and vaccine compositions thereof are useful for administration to mammals, including humans, to treat and/or prevent tumors associated with expression of MAGE antigen.

5 For pharmaceutical compositions, the peptides, i.e., the CTL or T helper peptides or CTL/T helper peptide conjugates, as described above will be administered to an individual already suffering from a MAGE-associated tumor. Those in the early stages of tumor development can be treated  
10 with the immunogenic peptides separately or in conjunction with other treatments, as appropriate. In therapeutic applications, compositions are administered to a patient in an amount sufficient to elicit an effective CTL response to a MAGE-bearing tumor and to cure or at least partially arrest  
15 its symptoms and/or complications. An amount adequate to accomplish this is defined as "therapeutically effective dose." Amounts effective for this use will depend on, e.g., the peptide composition, the manner of administration, the stage and severity of the disease being treated, the weight  
20 and general state of health of the patient, and the judgment of the prescribing physician, but generally range for the initial immunization (that is for therapeutic or prophylactic administration) from about 1.0  $\mu\text{g}$  to about 500  $\mu\text{g}$  of peptide for a 70 kg patient, followed by boosting dosages of from  
25 about 1.0  $\mu\text{g}$  to about 100  $\mu\text{g}$  of peptide pursuant to a boosting regimen over weeks to months depending upon the patient's response and condition by measuring specific CTL activity in the patient's blood. It must be kept in mind that the peptides and compositions of the present invention may  
30 generally be employed in serious disease states, that is, life-threatening or potentially life threatening situations. In such cases, in view of the minimization of extraneous substances and the relative nontoxic nature of the peptides, it is possible and may be felt desirable by the treating  
35 physician to administer substantial excesses of these peptide compositions.

Single or multiple administrations of the compositions can be carried out with dose levels and pattern

being selected by the treating physician. In any event, the pharmaceutical formulations should provide a quantity of CTL or T helper stimulatory peptides of the invention sufficient to effectively treat the patient.

5           As individuals may develop MAGE-associated tumors because of an inadequate (or absent) CTL or T helper response during the earliest stages of tumor development, it is important to provide an amount of immuno-potentiating peptide in a formulation and mode of administration sufficient to  
10           effectively stimulate a CTL or T helper cell response. Administration should continue until at least clinical symptoms or laboratory indicators indicate that the tumor has been eliminated or its progression has substantially abated and for a period thereafter. Immunizing doses followed by  
15           boosting doses at established intervals, e.g., from one to four weeks, may be required, possibly for a prolonged period of time, as necessary to resolve the tumor.

          The pharmaceutical compositions for therapeutic treatment are intended for parenteral, topical, oral or local  
20           administration. Preferably, the pharmaceutical compositions are administered parenterally, e.g., intravenously, subcutaneously, intradermally, or intramuscularly. Thus, the invention provides compositions for parenteral administration which comprise a solution of the CTL or T helper stimulatory  
25           peptides dissolved or suspended in an acceptable carrier, preferably an aqueous carrier. A variety of aqueous carriers may be used, e.g., water, buffered water, 0.4% saline, 0.3% glycine, hyaluronic acid and the like. These compositions may be sterilized by conventional, well known sterilization  
30           techniques, or may be sterile filtered. The resulting aqueous solutions may be packaged for use as is, or lyophilized, the lyophilized preparation being combined with a sterile solution prior to administration. The compositions may contain  
35           pharmaceutically acceptable auxiliary substances as required to approximate physiological conditions, such as pH adjusting and buffering agents, tonicity adjusting agents, wetting agents and the like, for example, sodium acetate, sodium

lactate, sodium chloride, potassium chloride, calcium chloride, sorbitan monolaurate, triethanolamine oleate, etc.

In some embodiments it may be desirable to include in the pharmaceutical composition at least one component which primes CTL. Certain lipids are capable of priming CTL responses in vivo, e.g., palmitic acid residues can be attached to the alpha and epsilon amino groups of a Lys residue and then linked to a synthetic peptide which comprises a class I-restricted CTL epitope. As further described herein, the lipidated peptide can then be incorporated into a liposome emulsified in an adjuvant, e.g., incomplete Freund's adjuvant. The arrangement of the components of the conjugate comprising the CTL inducing peptide/T helper peptide/lipid can be varied. In one case, the lipid moiety can be linked to the amino terminal end of the CTL inducing peptide, which in turn is linked at its carboxy terminal to the T helper peptide. In another case, the lipid is linked at the amino terminal end of the T helper peptide, which is linked at its carboxy terminal to the CTL inducing peptide. In each case, a spacer molecule, e.g., Lys-Ser-Ser, can be selectively inserted between the lipid moiety and the CTL or T helper peptide, as well as between the T helper and the CTL inducing peptides.

The concentration of CTL stimulatory peptides of the invention in the pharmaceutical formulations can vary widely, i.e., from less than about 1%, usually at or up to at least about 10%, to as much as 20 to 50% or more by weight, and will be selected primarily by fluid volumes, viscosities, etc., in accordance with the particular mode of administration selected. Thus, a typical pharmaceutical composition for intravenous infusion could be made up to contain 250 ml of sterile Ringer's solution, and 100 mg of peptide. Actual methods for preparing parenterally administrable compounds will be known or apparent to those skilled in the art and are described in more detail in for example, Remington's Pharmaceutical Sciences, 17th ed., Mack Publishing Company, Easton, PA (1985), which is incorporated herein by reference.

The peptides of the invention may also be administered via liposomes. Liposomes, which include

emulsions, foams, micelles, insoluble monolayers, phospholipid dispersions, lamellar layers and the like, can serve as a vehicle to target the peptides to a particular tissue, such as lymphoid tissue, or to tumor cells, as well as to increase the half-life of the peptide composition. In these preparations the peptide to be delivered is incorporated as part of a liposome, alone or in conjunction with a molecule which binds to, e.g., a receptor prevalent among lymphoid cells or tumor cells, such as monoclonal antibodies, or with other therapeutic or immunogenic compositions. A variety of methods are available for preparing liposomes, as described in, e.g., U.S. Patents 4,837,028 and 5,019,369, incorporated herein by reference.

In another aspect the present invention is directed to vaccines which contain as an active ingredient an immunogenically effective amount of a CTL stimulating peptide as described herein. The peptide(s) may be introduced into a host, including humans, linked to its own carrier or as a homopolymer or heteropolymer of active peptide units. Such a polymer has the advantage of increased immunological reaction and, where different peptides are used to make up the polymer, the additional ability to induce antibodies and/or cytotoxic T cells that react with different antigenic determinants of MAGE proteins. Useful carriers are well known in the art, and include, e.g., thyroglobulin, albumins such as human serum albumin, tetanus toxoid, polyamino acids such as poly(D-lysine:D-glutamic acid), influenza, hepatitis B virus core protein, and the like. The vaccines can also contain a physiologically tolerable (acceptable) diluent such as water, phosphate buffered saline, or saline, and further typically include an adjuvant. Adjuvants such as incomplete Freund's adjuvant, aluminum phosphate, aluminum hydroxide, or alum are materials well known in the art. Upon immunization with a peptide composition as described herein, via injection, aerosol, oral, transdermal or other route, the immune system of the host responds to the vaccine by producing large amounts of CTLs specific for MAGE antigen, and the host becomes at

least partially immune to MAGE-bearing tumors or resistant to such tumors.

Vaccine compositions containing the peptides of the invention are administered to a patient susceptible to or otherwise at risk of developing MAGE-associated tumors, such as melanomas, to enhance the patient's own immune response capabilities. Such an amount is defined to be a "immunogenically effective dose." In this use, the precise amounts again depend on the patient's state of health and weight, the mode of administration, the nature of the formulation, etc., but generally range from about 1.0  $\mu$ g to about 500  $\mu$ g per 70 kilogram patient, more commonly from about 50  $\mu$ g to about 100  $\mu$ g mg per 70 kg of body weight. CTL-inducing peptides are administered to individuals of an appropriate HLA type. In some instances it may be desirable to combine the peptide vaccines of the invention with vaccines which induce neutralizing cellular or antibody responses to tumor antigens, such as the p97 tumor antigen, for example.

For therapeutic or immunization purposes, the peptides of the invention can also be expressed by attenuated viral hosts, e.g., vaccinia or fowlpox. This approach involves the use of vaccinia virus as a vector to express nucleotide sequences that encode the peptides or conjugates of the invention. Upon introduction into a host, the recombinant vaccinia virus expresses the MAGE peptide, and thereby elicits a host CTL or T helper response to MAGE antigen on the tumor cells. Vaccinia vectors and methods useful in immunization protocols are described in, e.g., U.S. Patent No. 4,722,848, incorporated herein by reference. A wide variety of other vectors useful for therapeutic administration or immunization of the peptides of the invention will be apparent to those skilled in the art from the description herein.

The immunogenic peptides of the invention may be used to elicit CTL ex vivo. The resulting CTL can be used to treat tumors in patients that do not respond to other conventional forms of therapy, or will not respond to a peptide vaccine approach of therapy. Ex vivo CTL responses to a MAGE expressing tumor are induced by incubating in tissue

culture a patient's CTL precursor cells (CTLp) together with a source of antigen-presenting cells (APC) and the appropriate immunogenic peptide. After an appropriate incubation time (typically 1-4 weeks), in which the CTLp are activated and mature and expand into effector CTL, the cells are infused back into the patient, where they will inhibit or kill targeted tumor cells.

The peptides may also find use as diagnostic reagents. For example, a peptide of the invention may be used to determine the susceptibility of a particular individual to a treatment regimen which employs the peptide or related peptides, and thus may be helpful in modifying an existing treatment protocol or in determining a prognosis for an affected individual. In addition, the peptides may also be used to predict which individuals will be at substantial risk for developing MAGE-associated tumors.

The following examples are offered by way of illustration, not by way of limitation.

#### EXAMPLE 1

##### Cloning of Full Length MAGE DNA

The predicted coding region of MAGE-1 was cloned from first strand cDNA synthesized from the MAGE-1 positive cell line 938 by PCR amplification using primers located at bp605-622 and bp1459-1476. The amplification was carried out using a high fidelity thermostable polymerase, Pfu, by cycling 30 times, 2 minutes at 95°C, 2 minutes at 50°C, and 2 minutes at 73°C. These conditions resulted in the amplification of a 870 bp fragment consistent with the predicted size for MAGE-1. The amplified fragment was subcloned in the vector pRc/RSV (Invitrogen) for further characterization.

The cloned fragment was sequenced using sequenase (v 2.0, USB) and MAGE-1 specific primers. Sequence characterization of the cloned fragment found a cytidine inserted at nucleotide 1377. This insertion falls immediately 3' of the sequence.



To determine if the insertion had been introduced during PCR amplification the genomic sequence of MAGE-1 from the original cell line as well as from four independent normal individuals was determined. Genomic DNA was isolated and the MAGE-1 gene was amplified using the same conditions used in the original cloning. The amplified fragment was cycle sequenced using an antisense primer corresponding to nucleotides 1427-1448. The procedures were as follows.

Genomic DNA Isolation: Cells from the MAGE-1 expressing cell line were taken directly from tissue culture stocks. The procedure for genomic DNA isolation from individuals was as follows: 15 mls of heparinized whole blood was mixed with 15 mls of RPMI 1640 media and layered over 20 mls of lymphocyte separation media and centrifuged for 15 minutes at 400 x g following the manufacturers protocol (Ficoll-Paque, Pharmacia). The cell layer was collected and washed twice in RPMI 1640 media. The lymphocytes were counted, resuspended at  $4 \times 10^6$  cells/ml in 90% fetal bovine serum, 10% dimethyl sulfoxide and stored under liquid nitrogen until further processing.

The thawed cell pellet was lysed in 400  $\mu$ l of lysis buffer (4.2 M guanidine thiocyanate, 25.5 mM sodium acetate, 122 mM  $\beta$ -mercaptoethanol). The lysate was extracted once with an equal volume of phenol/chloroform and then an equal volume of chloroform. Sodium acetate was added to a final concentration of 0.3 M and the DNA was precipitated with two volumes of ethanol. The purified genomic DNA was resuspended in 200  $\mu$ l of H<sub>2</sub>O. The DNA concentration was determined by fluorometry following the directions provided by the manufacturer (TKO 0100 Fluorometer, Hoeffer).

#### DNA Amplification

A 100  $\mu$ l reaction mix was prepared containing 0.5  $\mu$ g of genomic DNA, 0.5  $\mu$ M of each amplification primer (primers indicated above), 10 mM Tris-HCl (pH 8.8), 50 mM KCl, 1.5 mM MgCl<sub>2</sub>, 0.001% (w/v) gelatin, all four deoxyribonucleotide triphosphates (dNTPs) each at 500  $\mu$ M, add 1.25 units of Taq DNA polymerase (Stratagene). The incubation conditions used

were: 30 cycles of 95°C, 2 minutes, 50°C, 2 minutes, 72°C, 2 minutes.

PCR DNA Fragment Purification:

The 100 µl reaction mix was fractionated by electrophoresis through a 1% agarose (SeaKem Agarose, FMC Inc.) gel, containing 10 µg/ml ethidium bromide, 40 mM Tris-acetate, 1 mM EDTA. A gel slice containing the desired DNA fragment (870 bp for the MAGE-1 amplified fragment) was excised during UV illumination. The DNA was purified using a glass bead purification kit (Qiaex, Qiagen) and was eluted in 20 µl of H<sub>2</sub>O.

DNA Sequencing:

All of the eluted DNA was sequenced with the appropriate primer (49 nucleotides downstream of the MAGE-1 amplified fragment) following the [<sup>35</sup>S]dCTP incorporation protocol recommended by the manufacturers of the commercial cycle-sequencing kit (ΔTaq Cycle-Sequencing Kit, Untied States Biochemical). The sequencing reactions were fractionated by electrophoresis through a 8% polyacrylamide gel (0.4 mm thick). The polyacrylamide gel was dried and exposed to X-ray film (XAR-5, Kodak) for 16 to 48 hours.

Each of the five MAGE 1 genes sequences contained the cytidine insertion at nucleotide 1377 when compared to the sequence in the GenBank entry. The insertion is significant because it shifts the reading frame of the gene, which changes 25 C-terminal amino acids and also extends the protein for an additional 33 amino acids. The full length DNA and amino acid sequence of human MAGE-1 protein is shown in Figure 1. Figure 2 shows the nucleotide and amino acid sequence of the newly discovered C terminal portion to the human MAGE-1 protein.

EXAMPLE 2Identification of MAGE Immunogenic Peptides  
from the C-Terminus

5           Using motifs for the various MHC class I alleles the C-terminus of the MAGE protein was analyzed for the presence of the motifs.

10           The motif for HLA-A3.2 comprises from the N-terminus to C-terminus a first conserved residue of L, M, I, V, S, A, T and F at position 2 and a second conserved residue of K, R or Y at the C-terminal end. Other first conserved residues are C, G or D and alternatively E. Other second conserved residues are H or F. The first and second conserved residues are preferably separated by 6 to 7 residues.

15           The motif for HLA-A1 comprises from the N-terminus to the C-terminus a first conserved residue of T, S or M, a second conserved residue of D or E, and a third conserved residue of Y. Other second conserved residues are A, S or T. The first and second conserved residues are adjacent and are preferably separated from the third conserved residue by 6 to 20 7 residues. A second motif consists of a first conserved residue of E or D and a second conserved residue of Y where the first and second conserved residues are separated by 5 to 6 residues.

25           The motif for HLA-A11 comprises from the N-terminus to the C-terminus a first conserved residue of T or V at position 2 and a C-terminal conserved residue of K. The first and second conserved residues are preferably separated by 6 or 7 residues.

30           The motif for HLA-A24.1 comprises from the N-terminus to the C-terminus a first conserved residue of Y, F or W at position 2 and a C terminal conserved residue of F, I, W, M or L. The first and second conserved residues are preferably separated by 6 to 7 residues.

35           The motif for HLA-A2.1 for 9 mer peptides includes one of the amino acids I, V, A and T at position L and V, L, I, A and M at position 9. Neither acidic amino acids nor P were found in position 1. Only one acidic amino acid and no

basic amino acids were found in position 3. Positions 6 and 7 showed no charged residues. Acidic amino acids, however, were frequently found in position 8, where they are tolerated, according to our definition of the A2.1 motif. The analysis of the sequences of naturally processed peptides therefore reveals that >90% of the peptides followed the defined rules for a complete motif.

The motif for HLA-A2.1 for 10 mer peptides includes one of the amino acids L, M, I, V, A and T at position 2 and V, I, L, A and M at position 10. In position 1 for example, in 10-mers again the P residue and acidic amino acids were not tolerated. In addition at position 1 in 10-mers aromatic residues were frequently observed in A2.1 binders. In position 3, acidic amino acids were frequently associated with poor binding capacity in both 9-mers and 10-mers. Interestingly, however, while in position 3 aromatic residues were preferred in 9-mers, aliphatic residues (L, V, I, M) were preferred in 10-mers.

The immunogenic peptides of about 9 and 10 amino acids in length identified using the motifs are set forth in Table 1.

Table 1

Antigen	Position	Sequence	Size	Allele
MAGE1N	274	TSYVKVLEY	9	A01
MAGE1N	254	VPDSDPARY	9	A01
MAGE1N	268	PRALAETSY	9	A01
MAGE1N	301	ALREEEEGV	9	A02
MAGE1N	264	FLWGPRALA	9	A02
MAGE1N	276	YVKVLEYVI	9	A02
MAGE1N	289	RVRFFFPSL	9	A02
MAGE1N	278	KVLEYVIKV	9	A02
MAGE1N	282	YVIKVSARV	9	A02
MAGE1N	269	RALAETSYV	9	A02
MAGE1N	271	LAETSYVKV	9	A02
MAGE1N	253	QVPDSDPAR	9	A03

	Antigen	Position	Sequence	Size	Allele
	MAGE1N	285	KVSARVRFF	9	A03
	MAGE1N	270	ALAETSYVK	9	A03
	MAGE1N	283	VIKVSARVR	9	A03
	MAGE1N	295	PSLREAAALR	9	A03
5	MAGE1N	274	TSYVKVLEY	9	A03
	MAGE1N	256	DSDPARYEF	9	A03
	MAGE1N	286	VSARVRFFF	9	A03
	MAGE1N	253	QVPDSDPAR	9	A11
	MAGE1N	283	VIKVSARVR	9	A11
10	MAGE1N	295	PSLREAAALR	9	A11
	MAGE1N	274	TSYVKVLEY	9	A11
	MAGE1N	270	ALAETSYVK	9	A11
	MAGE1N	244	VQEKYLEYR	9	A11
	MAGE1N	263	EFLWGPRAL	9	A24
15	MAGE1N	273	ETSYVKVLEY	10	A01
	MAGE1N	270	ALAETSYVKV	10	A02
	MAGE1N	279	VLEYVIKUSA	10	A02
	MAGE1N	271	LAETSYVKVL	10	A02
	MAGE1N	300	AALREEEEGV	10	A02
20	MAGE1N	276	YVKVLEYVIK	10	A03
	MAGE1N	243	LVQEKYLEYR	10	A03
	MAGE1N	282	YVIKVSARVR	10	A03
	MAGE1N	289	RVRFFFPSLR	10	A03
	MAGE1N	253	QVPDSDPARY	10	A03
25	MAGE1N	285	KVSARVRFFF	10	A03
	MAGE1N	242	DLVQEKYLEY	10	A03
	MAGE1N	283	VIKVSARVRF	10	A03
	MAGE1N	269	RALAETSYVK	10	A03
	MAGE1N	273	ETSYVKVLEY	10	A03
30	MAGE1N	255	PDSDPARYEF	10	A03
	MAGE1N	280	LEYVIKVSAR	10	A03
	MAGE1N	276	YVKVLEYVIK	10	A11
	MAGE1N	243	LVQEKYLEYR	10	A11

Antigen	Position	Sequence	Size	Allele
MAGE1N	282	YVIKVSARVR	10	A11
MAGE1N	289	RVRFFFP SLR	10	A11
MAGE1N	253	QVPDSDPARY	10	A11
MAGE1N	269	RALAETSYVK	10	A11
MAGE1N	273	ETSYVKVLEY	10	A11
MAGE1N	252	RQVPDSDPAR	10	A11
MAGE1N	280	LEYVIKVSAR	10	A11
MAGE1N	275	SYVKVLEYVI	10	A24
MAGE1N	293	FFPSLREAAAL	10	A24
MAGE1N	271	LAETSYVKVL	10	A24

These peptides were then evaluated as to their capacity to bind to the appropriate class I molecules using specific binding assays as described in copending and commonly owned application USSN 08/027,746, incorporated herein by reference. Results of the binding assays are set forth in table 2.

TABLE 2

## MAGE CTL TARGETS

Sequence	Posit.	Motif	Size	A1	A2	A3	A11	A24	Id
ETSYVKVLEY	274	A01/11	10	0.5600					1072.28
TSYVKVLEY	275	A01/11	9	0.0990					1072.29
KVLEYVIKV	279	A02	9		0.0900				1072.30
FLWGPRALA	265	A02	9		0.0420				1072.31
ALREEEGV	302	A02	9		0.0210				1072.32
ALAETSYVKV	271	A02	10		0.0150				1072.33
YVIKVSARV	283	A02	9		0.0140				1072.34
RALAETSYV	270	A02	9		0.0100				1072.35
TSYVKVLEY	275	A03/11	9			0.7100	0.0100		1072.36
RVRFFFP SLR	290	A03/11	10			0.4300	0.0089		1072.37
ALAETSYVK	271	A03/11	9			0.3100	0.3600		1072.38
RALAETSYVK	270	A03/11	10			0.1800	0.2400		1072.39
DLVQEKLEY	242	A03	10			0.0320	0.0051		1072.40
YVIKVSARVR	283	A03	10			0.0190	0.0009		1072.41
LVQEKLEY	243	A11	9			0.0026	0.0340		1072.42
SVYKVLEYVI	276	A24	10					0.0360	1072.43

Immunogenic peptides which were high and intermediate binders were then tested for their capacity to induce an *in vitro* CTL response.

5           The assay was carried out as follows:

To identify CTL epitopes, CTL was stimulated by SAC-I activated PBMCs as APC. Cold temperature expression of the MHC in which the  $\beta$ -2-microglobulin is unstable was utilized in addition to acid stripping to generate PBMC APC.

10           Complete Culture Medium. The tissue culture medium used in this study consisted of RPMI 1640 without Hepes (Biowhittaker) supplemented with 2 mM L-glutamine (Irvine Scientific), 0.5mM sodium pyruvate (Gibco), 100 U/100  $\mu$ g/ml penicillin/ streptomycin (Irvine), and 5% heat-inactivated  
15 Human Serum Type AB (RPMI/5% HS; Gemini Bioproducts). Culture media used in the growth of EBV-transformed lines contained 10% heat-inactivated fetal calf serum (RPMI/10% FCS, Irvine) instead of human serum.

Cytokines. Recombinant human interleukin-2 (rIL-2) and interleukin-4 (rIL-4) were obtained from Sandoz and used  
20 at a final concentration of 10  $\mu$ g/ml and 10  $\mu$ g/ml, respectively. Human interferon- $\gamma$  (IFN- $\gamma$ ) and recombinant human Interleukin-7 (rIL-7) were obtained from Genzyme and used at 20 U/ml and 10 ng/ml, respectively.

25           Peptides. Peptides were synthesized on an automated synthesizer and are described in Table 1. Peptides were routinely diluted in 100% DMSO at 20 mg/ml, aliquoted, and stored at -70°C. Pools of peptides were tested with 2-3  
30 peptides/pool with no more than 5 fold differences in Class I binding (if a pool was not possible for a specific peptide then individual peptides were tested).

Cell Lines. JY, Steinlin, EHM, BVR, and KT3 are homozygous human EBV-transformed B cell lines expressing HLA  
A<sub>2.1</sub>, A<sub>1</sub>, A<sub>3</sub>, A<sub>11</sub>, and A<sub>24</sub>, respectively. They are grown in  
35 RPMI/10% FCS. K562, an NK cell sensitive, erythoblastoma line grown in RPMI/10% FCS, was used for reduction of background killing.



Isolation of Peripheral Blood Mononuclear Cells (PBMCs). Whole blood was collected into heparin containing syringes and spun in 50cc tubes at 1600 RPM (Beckman GS-6KR) for 15 minutes. The plasma layer was then removed and 10 ml of buffy coat was collected with a pipette using a circular motion (an additional 2 ml from the bottom of the tube was included in the 10 ml). The buffy coat was mixed well and diluted with an equal volume of RPMI. The buffy coat (30 ml) was then layered on 20 ml of Ficoll-Paque (Pharmacia) and centrifuged at 1850 RPM (400xg) for 20 minutes, 25°C, with the brake off. The interface between the ficoll and the plasma containing the PBMCs was recovered with a transfer pipet (two interfaces per 50 ml tube) and washed three times with 50 ml of RPMI (1700, 1500, and 1300 RPM for 10 minutes). Cells were resuspended in 10-20 ml of culture medium, counted, and adjusted to the appropriate concentration.

Freezing PBMCs. 30 million cells/tube (90% FCS/10% DMSO; Sigma) were inserted into a Nalgene Cryo 1°C Freezing Container containing Isopropanol (Fisher) and placed at -70°C from 4 hrs (minimum) to overnight (maximum). The Isopropanol was changed every five times. Tubes were transferred to liquid nitrogen for long term storage. To thaw, PBMCs were continuously shaken in a 37°C water bath until the last crystal was almost thawed (tubes were not allowed to sit in the water bath or at room temperature for any period of time). Cells were diluted into serum-free RPMI and washed twice.

#### Induction of Primary CTL Using SAC-I Activated PBMCs as APCs

a. Preparation of APCs: PBMCs were purified using the standard Ficoll-Paque protocol and resuspended at  $1 \times 10^6$ /ml in RPMI/5% FCS containing 0.005% Pansorbin cells (SAC-I cells expressing Protein A; Calbiochem), 20 µg/ml Immunobeads (Rabbit anti-Human IgM; Biorad), and 20 ng/ml of human rIL-4. Two ml of cells per well were plated in a 24-well plate (Falcon, Becton Dickinson) and cultured at 37°C. After 3 days, the medium was removed and the cells were washed three times followed by addition of RPMI/10% HS. The cells were used after culturing for an additional 2 days in RPMI/10% HS.

b. Expression of empty Class I molecules on the surface of APCs and peptide loading of APCs.

1. Cold temperature incubation:

a. Expression of empty MHC in APCs: The APCs were adjusted to a concentration of  $2 \times 10^6$ /ml in complete culture medium (Section #1) containing 10 ng/ml rIL-4, 20 U/ml human IFN- $\gamma$ , and 3  $\mu$ g/ml  $\beta_2$ -microglobulin ( $\beta_2$ m). The cells were then incubated overnight at 26°C in the presence of 5% CO<sub>2</sub>. These cells express only a fraction of Class I molecules in the empty state (~10%).

b. Peptide loading of APC stimulator cells:

Empty Class I expressing APCs were washed 1-2 times with serum free RPMI (+ L-glutamine and Hepes) and resuspended at  $1 \times 10^7$  in serum-free RPMI containing 50  $\mu$ g/ml total of the peptide pool (i.e., 16.7  $\mu$ g/ml of each peptide in a pool of three; 25  $\mu$ g/ml of each peptide in a pool of two; 50  $\mu$ g/ml of individual peptide), 30  $\mu$ g/ml DNase, and 3  $\mu$ g/ml  $\beta_2$ m. Following a 4 hour incubation at 20°C, the cells were irradiated at 6100 rads ( $5 \times 10^6$ / ml; 25 million cells/tube), washed and adjusted to the appropriate concentration for addition to the induction culture (see below).

2. Acid stripping: This was used as an

alternative method for generating empty MHC on the surface of the APCs. The SAC-I activated PBMCs were washed once in cold 0.9% sodium chloride (J.T. Baker) containing 1% BSA. The cells were resuspended at  $10^7$ /ml in cold citrate-phosphate buffer (0.13M L-ascorbic acid [J.T. Baker], 0.06M sodium phosphate monobasic [Sigma], pH3) containing 1% BSA and 3  $\mu$ g/ml  $\beta_2$ m and incubated on ice. After 2 minutes, 5 volumes of cold 0.15M sodium phosphate monobasic buffer, pH7.5, containing 1% BSA, 3  $\mu$ g/ml  $\beta_2$ m, and 10  $\mu$ g/ml peptide [neutralizing buffer #1] was added and the cells centrifuged at 1500 RPM for 5 minutes at 4°C. The cells were resuspended in 1 ml of cold PBS containing 1% BSA, 30  $\mu$ g/ml DNase, 3  $\mu$ g/ml  $\beta_2$ -microglobulin, and 50  $\mu$ g/ml peptide [neutralizing buffer #2] and incubated for 4 hours at 20°C. As above, subsequent to the four hour incubation at 20°C, the cells were irradiated at 6100 rads ( $5 \times 10^6$ / ml; 25 million cells/tube), washed, then

adjusted to the appropriate concentration for addition to the induction culture (see below).

5           c.   Preparation of the CD4+ depleted PBMC responder cell population (depletion of lymphocyte sub-populations using AIS flasks). AIS MicroCollector T-150 flasks (specific for the depletion of CD4+ T cells; Menlo Park, CA) were primed by adding 25 ml of PBS/1 mM EDTA, swirling for 30 seconds so that all surfaces were moistened, and then  
10           incubating with the binding surface down at room temperature for 1 hour. Following this incubation, flasks were shaken vigorously for 30 seconds, washed 1 time with PBS/EDTA, 2 additional times with PBS and then incubated with 25 ml of culture medium for 15 minutes. PBMCs were thawed in  
15           serum-free RPMI (+ L-glutamine + Hepes) containing 30  $\mu$ g/ml DNase, washed once, and incubated for 15 minutes in culture medium. Following aspiration of culture medium from the flasks, up to 180 million PBMCs were added in 25 ml of culture medium containing 30  $\mu$ g/ml DNase. After 1 hour at room  
20           temperature, the flasks were rocked gently for 10 seconds to resuspend the nonadherent cells. The nonadherent cell suspension containing the CD8+ T cells was collected and the flasks were washed 2 times with PBS. The CD4+ T cell depleted PBMCs were centrifuged and counted for addition to the  
25           induction culture. The CD4+ and CD8+ phenotype of the CD4+ depleted cell population was determined by FACS analysis (see below). In general, this technique resulted in a two-fold enrichment for CD8+ T cells with an average of approximately 40-50% CD8+ T cells and 15-20% remaining CD4+ T cells  
30           following depletion of CD4+ T cells.

          d.   Induction of primary CTL. During the 4 hour peptide loading of the stimulator APCs, CD4+ depleted PBMC to be used as the responder population were prepared utilizing AIS flasks for selection of CD8+ T cells through the depletion of CD4+ T  
35           cells (above). The responder cells were plated at  $3 \times 10^6$ /ml in a 1 ml volume (24 well plate) and placed at 37°C until the peptide loaded stimulator APCs were prepared. The irradiated, peptide loaded APCs were washed 1 time in serum-free RPMI (+

L-glutamine and Hepes), adjusted to the appropriate concentration in complete medium, and plated into a 24 well plate at 1 ml/plate: For PBMC,  $1 \times 10^6$  stimulator cells (1 ml volume) were plated into the wells containing the responder cells; For SAC-I activated PBMC and PHA blasts, 1 ml of  $3 \times 10^5$ /ml stimulator cells were plated in each well. A final concentration of 10  $\mu$ g/ml of additional peptide was added in addition to 10 ng/ml final concentration of rIL-7 (2 ml total volume). The cells were cultured for 12 days. (For the "pulse only" induction protocol, the additional 10  $\mu$ g/ml of soluble peptide was not added to the cultures). On day 12, the cultures were restimulated with peptide pulsed adherent cells and tested for cytolytic activity 7 days later (below).

Protocol for Restimulation of Primary CTL Using Adherent APC. PBMCs were thawed into serum-free RPMI (+ L-glutamine and Hepes) containing 30  $\mu$ g/ml DNase, washed 2 times, and adjusted to  $5 \times 10^6$  /ml in culture medium containing DNase. PBMCs (25 million cells/tube in 5 ml) were irradiated at 6100R. After 1 wash, the PBMCs were resuspended in culture medium and adjusted to  $4 \times 10^6$ /ml. 1 ml of irradiated PBMCs was added per well of a 24-well plate. The PBMC were incubated for 2 hours at 37°C, washed 3 times to remove non-adherent cells, and cultured in medium containing 20  $\mu$ g/ml total peptide and 3  $\mu$ g/ml  $\beta_2$ microglobulin added in a 0.5 ml volume and again incubated for 2 hours at 37°C. The peptide was aspirated and  $1.5 \times 10^6$  responder cells resuspended in culture medium were added in a 1 ml volume. After 2 days, 1 ml of culture medium containing 20 U/ml rIL-2 was added.

FACS Analysis. One million cells/tube were centrifuged, resuspended in 100  $\mu$ l/tube PBS/0.1% BSA/0.02% sodium azide (Sigma) plus 10  $\mu$ l/tube directly conjugated antibody (Becton Dickinson), and incubated on ice 15-20 minutes. Cells were then washed 2 times with PBS/0.1% BSA/0.02% sodium azide and resuspended in PBS to analyze on FACScan (Becton Dickinson). When it was not possible to analyze samples within 1-2 days, cells were fixed with PBS

containing 1% paraformaldehyde (Fisher) and analyzed within one week.

#### Cytotoxicity Assays

a. Target cell preparation. Approximately 16-20 hours prior to the CTL assay, target cells (Class I matched EBV-transformed lines) were washed once and resuspended in a 10 ml volume at  $3 \times 10^5$ /ml in RPMI/5% FCS in the presence or absence of 10  $\mu$ g/ml total peptide.

b. Labeling of target cells: Target cells were centrifuged and resuspended in 200  $\mu$ l/tube sodium  $^{51}\text{Cr}$  chromate (NEN), then incubated at 37°C for 1 hour on a shaker. Targets were washed 3 times (10 ml/wash) with RPMI/10% FCS and resuspended in 10 ml (to determine the efficiency of labelling, 50  $\mu$ l/target was counted on the Micromedic automatic gamma counter).

c. CTL assay. Target cells were adjusted to  $2 \times 10^5$ /ml and 50  $\mu$ l of the cell culture was added to each well of a U-bottomed 96-well plate (Costar Corp.) for a final concentration of  $1 \times 10^4$ /well. K562 cells were washed once, resuspended at  $4 \times 10^6$ /ml, and 50  $\mu$ l/well was added for a final concentration of  $2 \times 10^5$ /well (ratio of cold K562 to target was 20:1). Responder cells were washed once, resuspended at  $9 \times 10^6$ /ml, and three fold serial dilutions were performed for effector to target ratios of 90:1, 30:1, 10:1, and 3:1. Responder cells were added in a volume of 100  $\mu$ l in duplicate wells. For spontaneous release, 50  $\mu$ l/well of labelled target cells, 50  $\mu$ l/well K562, and 100  $\mu$ l/well of medium was added. For maximum release, 50  $\mu$ l/well target, 50  $\mu$ l/well K562, and 100  $\mu$ l/well of 0.1% Triton-X100 (Sigma) was added. Plates were centrifuged for 5 minutes at 1200 RPM. Following a 5 hour incubation at 37°C, plates were centrifuged again for 5 minutes at 1200 RPM, and 100  $\mu$ l/well of supernatant was collected. Standard gamma counting techniques (Micromedic automatic gamma counter; 0.5 minutes/tube) were used to determine the percent specific lysis according to the formula: % specific lysis =  $\frac{\text{cpm experimental} - \text{cpm spontaneous release}}{\text{cpm maximum release} - \text{cpm spontaneous release}} \times 100$ . A CTL assay was considered positive if the lysis by CTL of

targets sensitized with a specific peptide at the two highest effector to target (E:T) ratios was 15% greater than lysis of control targets (i.e., target cells without peptide). A cytotoxicity assay was considered borderline if the lysis by CTL of targets sensitized with a specific peptide at the two highest E:T ratios was 6% greater than lysis of control targets. Of the 63 MAGE peptides that bind to the indicated alleles 12 induced primary CTL responses.

Results of the CTL assays are shown in Fig. 3.

Although the present invention has been described in some detail by way of illustration and example for purposes of clarity and understanding, it will be apparent that certain changes and modifications may be practiced within the scope of the appended claims.

## SEQUENCE LISTING

## (1) GENERAL INFORMATION:

## (i) APPLICANT:

(A) NAME: Cytel Corporation  
 (B) STREET: 3525 John Hopkins Court  
 (C) CITY: San Diego  
 (D) STATE: California  
 (E) COUNTRY: United States of America  
 (F) POSTAL CODE (ZIP): 92121  
 (G) TELEPHONE: (619) 552-3000  
 (H) TELEFAX: (619) 552-8801  
 (I) TELEX:

(ii) TITLE OF INVENTION: CLONING AND CHARACTERIZATION OF THE  
 COMPLETE MAGE 1 GENE

(iii) NUMBER OF SEQUENCES: 24

## (iv) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: Floppy disk  
 (B) COMPUTER: IBM PC compatible  
 (C) OPERATING SYSTEM: PC-DOS/MS-DOS  
 (D) SOFTWARE: PatentIn Release #1.0, Version #1.25

## (v) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER: WO  
 (B) FILING DATE:  
 (C) CLASSIFICATION:

## (vi) PRIOR APPLICATION DATA:

(A) APPLICATION NUMBER: US 08/103,623  
 (B) FILING DATE: 06-AUG-1993

## (vii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Parmelee, Steven W.  
 (B) REGISTRATION NUMBER: 31,990  
 (C) REFERENCE/DOCKET NUMBER: 14137-60PC

## (viii) TELECOMMUNICATION INFORMATION:

(A) TELEPHONE: (206) 467-9600  
 (B) TELEFAX: (415) 543-5043

## (2) INFORMATION FOR SEQ ID NO:1:

## (i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 58 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

Arg	Gln	Val	Pro	Asp	Ser	Asp	Pro	Ala	Arg	Tyr	Glu	Phe	Leu	Trp	Gly
1					5				10					15	
Pro	Arg	Ala	Leu	Ala	Glu	Thr	Ser	Tyr	Val	Lys	Val	Leu	Glu	Tyr	Val
		20					25					30			
Ile	Lys	Val	Ser	Ala	Arg	Val	Arg	Phe	Phe	Phe	Pro	Ser	Leu	Arg	Glu
		35					40					45			

Ala Ala Leu Arg Glu Glu Glu Glu Gly Val  
50 55

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 10 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

Glu Thr Ser Tyr Val Lys Val Leu Glu Tyr  
1 5 10

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 9 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

Thr Ser Tyr Val Lys Val Leu Glu Tyr  
1 5

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 9 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

Lys Val Leu Glu Tyr Val Ile Lys Val  
1 5

(2) INFORMATION FOR SEQ ID NO:5:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 9 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

Phe Leu Trp Gly Pro Arg Ala Leu Ala



1 5

(2) INFORMATION FOR SEQ ID NO:6:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 9 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

Ala Leu Arg Glu Glu Glu Gly Val  
1 5

(2) INFORMATION FOR SEQ ID NO:7:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 10 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

Ala Leu Ala Glu Thr Ser Tyr Val Lys Val  
1 5 10

(2) INFORMATION FOR SEQ ID NO:8:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 9 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

Tyr Val Ile Lys Val Ser Ala Arg Val  
1 5

(2) INFORMATION FOR SEQ ID NO:9:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 11 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

Arg Ala Leu Ala Glu Thr Ser Tyr Val Lys Val  
1 5 10

## (2) INFORMATION FOR SEQ ID NO:10:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 9 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

Thr Ser Tyr Val Lys Val Leu Glu Tyr  
1 5

## (2) INFORMATION FOR SEQ ID NO:11:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 10 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

Arg Val Arg Phe Phe Phe Pro Ser Leu Arg  
1 5 10

## (2) INFORMATION FOR SEQ ID NO:12:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 11 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

Ala Leu Ala Glu Thr Ser Tyr Val Lys Val Lys  
1 5 10

## (2) INFORMATION FOR SEQ ID NO:13:

- (i) SEQUENCE CHARACTERISTICS:  
(A) LENGTH: 12 amino acids  
(B) TYPE: amino acid  
(C) STRANDEDNESS: single  
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

Arg Ala Leu Ala Glu Thr Ser Tyr Val Lys Val Lys  
1 5 10

## (2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 10 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

Asp Leu Val Gln Glu Lys Tyr Leu Glu Tyr  
1 5 10

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 10 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

Tyr Val Ile Lys Val Ser Ala Arg Val Arg  
1 5 10

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 9 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

Leu Val Gln Glu Lys Tyr Leu Glu Tyr  
1 5

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 10 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

Ser Tyr Val Lys Val Leu Glu Tyr Val Ile  
1 5 10

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 14 amino acids
- (B) TYPE: amino acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

Gln Tyr Ile Lys Ala Asn Ser Lys Phe Ile Gly Ile Thr Glu  
1                      5                      10

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 16 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

Lys Ile Ala Lys Met Lys Ala Ser Ser Val Phe Asn Val Val Asn Ser  
1                      5                      10                      15

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 20 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

Asp Ile Glu Lys Lys Ile Ala Lys Met Lys Ala Ser Ser Val Phe Asn  
1                      5                      10                      15

Val Val Asn Ser  
20

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
  - (A) LENGTH: 14 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

Ile Ser Gln Ala Val His Ala Ala His Ala Glu Ile Asn Glu  
1                      5                      10

## (2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 13 amino acids  
 (B) TYPE: amino acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

Pro Lys Tyr Val Lys Gln Asn Thr Leu Lys Leu Ala Thr  
 1 5 10

## (2) INFORMATION FOR SEQ ID NO:23:

- (i) SEQUENCE CHARACTERISTICS:  
 (A) LENGTH: 2420 base pairs  
 (B) TYPE: nucleic acid  
 (C) STRANDEDNESS: single  
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

## (ix) FEATURE:

- (A) NAME/KEY: CDS  
 (B) LOCATION: 626..1552

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

```

GGATCCAGGC CCTGCCAGGA AAAATATAAG GGCCCTGCGT GAGAACAGAG GGGGTCATCC      60
ACTGCATGAG AGTGGGGATG TCACAGAGTC CAGCCCACCC TCCTGGTAGC ACTGAGAAGC      120
CAGGGCTGTG CTTGCGGTCT GCACCCTGAG GGCCCGTGGA TTCCTCTTCC TGGAGCTCCA      180
GGAACCAGGC AGTGAGGCCT TGGTCTGAGA CAGTATCCTC AGGTCACAGA GCAGAGGATG      240
CACAGGGTGT GCCAGCAGTG AATGTTTGCC CTGAATGCAC ACCAAGGGCC CCACCTGCCA      300
CAGGACACAT AGGACTCCAC AGAGTCTGGC CTCACCTCCC TACTGTCAGT CCTGTAGAAT      360
CGACCTCTGC TGGCCGGCTG TACCCTGAGT ACCCTCTCAC TTCCTCCTTC AGGTTTTTCAG      420
GGGACAGGCC AACCAGAGG ACAGGATTCC CTGGAGGCCA CAGAGGAGCA CCAAGGAGAA      480
GATCTGTAAG TAGGCCTTTG TTAGAGTCTC CAAGGTTTCAG TTCTCAGCTG AGGCCTCTCA      540
CACACTCCCT CTCTCCCCAG GCCTGTGGGT CTTCATTGCC CAGCTCCTGC CCACACTCCT      600
GCCTGCTGCC CTGACGAGAG TCATCATGTC TCTTGAGCAG AGGAGTCTGC ACTGCAAGCC      660
TGAGGAAGCC CTTGAGGCCC AACAAAGAGC CCTGGGCCTG GTGTGTGTGC AGGCTGCCAC      720
CTCCTCCTCC TCTCCTCTGG TCCTGGGCAC CCTGGAGGAG GTGCCCACTG CTGGGTCAAC      780
AGATCCTCCC CAGAGTCCTC AGGGAGCCTC CGCCTTTCCC ACTACCATCA ACTTCACTCG      840
ACAGAGGCAA CCCAGTGAGG GTTCCAGCAG CCGTGAAGAG GAGGGGCCAA GCACCTCTTG      900
TATCCTGGAG TCCTTGTTCC GAGCAGTAAT CACTAAGAAG GTGGCTGATT TGGTTGGTTT      960
TCTGCTCCTC AAATATCGAG CCAGGGAGCC AGTCACAAAG GCAGAAATGC TGGAGAGTGT     1020

```

```

CATCAAAAAT TACAAGCACT GTTTCCTGA GATCTTCGGC AAAGCCTCTG AGTCCTTGCA 1080
GCTGGTCTTT GGCATTGACG TGAAGGAAGC AGACCCACC GGCCACTCCT ATGTCCTTGT 1140
CACCTGCCTA GGTCTCTCCT ATGATGGCCT GCTGGGTGAT AATCAGATCA TGCCCAAGAC 1200
AGGCTTCCTG ATAATTGTCC TGGTCATGAT TGCAATGGAG GGCGGCCATG CTCCTGAGGA 1260
GGAAATCTGG GAGGAGCTGA GTGTGATGGA GGTGTATGAT GGGAGGGAGC ACAGTGCCTA 1320
TGGGGAGCCC AGGAAGCTGC TCACCCAAGA TTTGGTGACG GAAAAGTACC TGGAGTACCG 1380
GCAGGTGCCG GACAGTGATC CCGCACGCTA TGAGTTCCTG TGGGGTCCAA GGGCCCTCGC 1440
TGAAACCAGC TATGTGAAAG TCCTTGAGTA TGTGATCAAG GTCAGTGCAA GAGTTCGCTT 1500
TTTCTTCCCA TCCCTGCGTG AAGCAGCTTT GAGAGAGGAG GAAGAGGGAG TCTGAGCATG 1560
AGTTGCAGCC AAGGCCAGTG GGAGGGGGAC TGGGCCAGTG CACCTTCCAG GGCCGCGTCC 1620
AGCAGCTTCC CCTGCCTCGT GTGACATGAG GCCCATTCTT CACTCTGAAG AGAGCGGTCA 1680
GTGTTCTCAG TAGTAGGTTT CTGTTCTATT GGGTGACTTG GAGATTATC TTTGTTCTCT 1740
TTTGGAATTG TTCAAATGTT TTTTTTTAAG GGATGGTTGA ATGAACTTCA GCATCCAAGT 1800
TTATGAATGA CAGCAGTCAC ACAGTTCTGT GTATATAGTT TAAGGGTAAG AGTCTTGTGT 1860
TTTATTGAGA TTGGGAAATC CATTCTATTT TGTGAATTGG GATAATAACA GCAGTGGAAAT 1920
AAGTACTTAG AAATGTGAAA AATGAGCAGT AAAATAGATG AGATAAAGAA CTAAAGAAAT 1980
TAAGAGATAG TCAATTCTTG CCTTATACCT CAGTCTATTC TGTAATTTT TAAAGATAT 2040
ATGCATACCT GGATTTCTT GGCTTCTTTG AGAATGTAAG AGAAATTAAA TCTGAATAAA 2100
GAAITCTTCC TGTTCACTGG CTCTTTTCTT CTCCATGCAC TGAGCATCTG CTTTTTGGAA 2160
GGCCCTGGGT TAGTAGTGGA GATGCTAAGG TAAGCCAGAC TCATACCCAC CCATAGGGTC 2220
GTAGAGTCTA GGAGCTGCAG TCACGTAATC GAGGTGGCAA GATGTCCTCT AAAGATGTAG 2280
GGAAAAGTGA GAGAGGGGTG AGGGTGTGGG GCTCCGGGTG AGAGTGGTGG AGTGTCAATG 2340
CCCTGAGCTG GGGCATTTTG GGCTTTGGGA AACTGCAGTT CCTTCTGGGG GAGCTGATTG 2400
TAATGATCTT GGGTGGATCC 2420

```

## (2) INFORMATION FOR SEQ ID NO:24:

- (i) SEQUENCE CHARACTERISTICS:
- (A) LENGTH: 309 amino acids
  - (B) TYPE: amino acid
  - (C) STRANDEDNESS: single
  - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: protein

## (xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

```

Met Ser Leu Glu Gln Arg Ser Leu His Cys Lys Pro Glu Glu Ala Leu
1           5           10           15
Glu Ala Gln Gln Glu Ala Leu Gly Leu Val Cys Val Gln Ala Ala Thr
20           25           30
Ser Ser Ser Ser Pro Leu Val Leu Gly Thr Leu Glu Glu Val Pro Thr

```

35					40					45					
Ala	Gly	Ser	Thr	Asp	Pro	Pro	Gln	Ser	Pro	Gln	Gly	Ala	Ser	Ala	Phe
50						55					60				
Pro	Thr	Thr	Ile	Asn	Phe	Thr	Arg	Gln	Arg	Gln	Pro	Ser	Glu	Gly	Ser
65					70					75					80
Ser	Ser	Arg	Glu	Glu	Glu	Gly	Pro	Ser	Thr	Ser	Cys	Ile	Leu	Glu	Ser
				85					90					95	
Leu	Phe	Arg	Ala	Val	Ile	Thr	Lys	Lys	Val	Ala	Asp	Leu	Val	Gly	Phe
			100					105					110		
Leu	Leu	Leu	Lys	Tyr	Arg	Ala	Arg	Glu	Pro	Val	Thr	Lys	Ala	Glu	Met
		115					120					125			
Leu	Glu	Ser	Val	Ile	Lys	Asn	Tyr	Lys	His	Cys	Phe	Pro	Glu	Ile	Phe
	130					135					140				
Gly	Lys	Ala	Ser	Glu	Ser	Leu	Gln	Leu	Val	Phe	Gly	Ile	Asp	Val	Lys
145				150						155					160
Glu	Ala	Asp	Pro	Thr	Gly	His	Ser	Tyr	Val	Leu	Val	Thr	Cys	Leu	Gly
				165					170					175	
Leu	Ser	Tyr	Asp	Gly	Leu	Leu	Gly	Asp	Asn	Gln	Ile	Met	Pro	Lys	Thr
			180					185					190		
Gly	Phe	Leu	Ile	Ile	Val	Leu	Val	Met	Ile	Ala	Met	Glu	Gly	Gly	His
		195					200					205			
Ala	Pro	Glu	Glu	Glu	Ile	Trp	Glu	Glu	Leu	Ser	Val	Met	Glu	Val	Tyr
	210					215					220				
Asp	Gly	Arg	Glu	His	Ser	Ala	Tyr	Gly	Glu	Pro	Arg	Lys	Leu	Leu	Thr
225				230						235					240
Gln	Asp	Leu	Val	Gln	Glu	Lys	Tyr	Leu	Glu	Tyr	Arg	Gln	Val	Pro	Asp
				245					250					255	
Ser	Asp	Pro	Ala	Arg	Tyr	Glu	Phe	Leu	Trp	Gly	Pro	Arg	Ala	Leu	Ala
			260					265					270		
Glu	Thr	Ser	Tyr	Val	Lys	Val	Leu	Glu	Tyr	Val	Ile	Lys	Val	Ser	Ala
		275					280					285			
Arg	Val	Arg	Phe	Phe	Phe	Pro	Ser	Leu	Arg	Glu	Ala	Ala	Leu	Arg	Glu
	290					295					300				
Glu	Glu	Glu	Gly	Val											
305															

What is claimed:

1. An immunogenic peptide comprising about nine to  
5 ten amino acids, wherein said peptide is from the C terminal  
58 amino acids of Sequence ID. No. 1.

2. The immunogenic peptide of claim 1 having an  
HLA-A 1/11 binding motif wherein the immunogenic peptide is  
10 selected from the group consisting of:

274MAGE1N Glu-Thr-Ser-Tyr-Val-Lys-Val-Leu-Glu-Tyr,  
(A01/11) [Seq. ID No. 2]; and  
275 MAGE1N Thr-Ser-Tyr-Val-Lys-Val-Leu-Glu-Tyr,  
(A01/11) [Seq. ID No. 3];

3. The immunogenic peptide of claim 1 having an  
HLA-A2 binding motif wherein the immunogenic peptide is  
selected from the group consisting of:

279 MAGE1N Lys-Val-Leu-Glu-Tyr-Val-Ile-Lys-Val,  
20 (A02) [Seq. ID No. 4];  
265 MAGE1N Phe-Leu-Trp-Gly-Pro-Arg-Ala-Leu-Ala,  
(A02) [Seq. ID No. 5];  
302 MAGE1N Ala-Leu-Arg-Glu-Glu-Glu-Glu-Gly-Val,  
A02 [Seq. ID No. 6];  
25 271 MAGE1N Ala-Leu-Ala-Glu-Thr-Ser-Tyr-Val-Lys-Val,  
A02 [Seq. ID No. 7];  
283 MAGE1N Tyr-Val-Ile-Lys-Val-Ser-Ala-Arg-Val,  
(A02) [Seq. ID No. 8]; and  
270 MAGE1N Arg-Ala-Leu-Ala-Glu-Thr-Ser-Tyr-Val,  
30 (A02) [Seq. ID No. 9];

4. The immunogenic peptide of claim 1 having an HLA  
A3/A11 binding motif wherein the immunogenic peptide is  
selected from the group consisting of:

35 275 MAGE1N Thr-Ser-Tyr-Val-Lys-Val-Leu-Glu-Tyr,  
(AA03/11) [Seq. ID No. 10];  
290 MAGE1N Arg-Val-Arg-Phe-Phe-Phe-Pro-Ser-Leu-Arg,  
(A03/11) [Seq. ID No. 11];



271 MAGE1N Ala-Leu-Ala-Glu-Thr-Ser-Tyr-Val-Lys-Val-Lys,  
A03/11 [Seq. ID No. 12];

270 MAGE1N Arg-Ala-Leu-Ala-Glu-Thr-Ser-Tyr-Val-Lys,  
(A03/11) [Seq. ID No. 13];

5 242 MAGE1N Asp-Leu-Val-Gln-Glu-Lys-Tyr-Leu-Glu-Tyr,  
(A03) [Seq. ID No. 14];

283 MAGE1N Tyr-Val-Ile-Lys-Val-Ser-Ala-Arg-Val-Arg,  
(A03) [Seq. ID No. 15]; or

10 243 MAGE1N Leu-Val-Gln-Glu-Lys-Tyr-Leu-Glu-Tyr,  
(A11) [Seq. ID No. 16];

5. The immunogenic peptide of claim 1 having an  
HLA-A24 binding motif wherein the immunogenic peptide is:

15 276 MAGE1N Ser-Tyr-Val-Lys-Val-Leu-Glu-Tyr-Val-Ile,  
(A24) [Seq. ID No. 17].

6. An isolated DNA encoding an immunogenic peptide  
selected from the C terminal 58 amino acids of Sequence I.D.  
No. 1.

20

7. A vector comprising DNA encoding an immunogenic  
peptide selected from the C terminal 58 amino acids of  
Sequence I.D. No. 1.

25

8. A host cell transformed with the vector of claim  
6.

Figure 1 (Sheet 1 of 3)

Sequence Range: 1 to 2420

10	20	30	40	50	60
GGATC	CAGGC	CCTGC	CAGGA	AAAAT	ATAAG
70	80	90	100	110	120
ACTGC	ATGAG	AGTGG	GGATG	TCACA	GAGTC
130	140	150	160	170	180
CAGGG	CTGTG	CTTGC	GGTCT	GCACC	CTGAG
190	200	210	220	230	240
GGAAC	CAGGC	AGTGA	GGCCT	TGGTC	TGAGA
250	260	270	280	290	300
CACAG	GGTGT	GCCAG	CAGTG	AATGT	TTGCC
310	320	330	340	350	360
CAGGA	CACAT	AGGAC	TCCAC	AGAGT	CTGGC
370	380	390	400	410	420
CGACC	TCTGC	TGGCC	GGCTG	TACCC	TGAGT
430	440	450	460	470	480
GGGAC	AGGCC	AACCC	AGAGG	ACAGG	ATTCC
490	500	510	520	530	540
GATCT	GTAAG	TAGGC	CTTTG	TTAGA	GTCTC
550	560	570	580	590	600
CACAC	TCCCT	CTCTC	CCCAG	GCCTG	TGGGT
610	620	630	640	650	
GCCTG	CTGCC	CTGAC	GAGAG	TCATC	ATG
					TCT
					CTT
					GAG
					CAG
					AGG
					AGT
					CTG
					CAC
					TGC
					Met
					Ser
					Leu
					Glu
					Gln
					Arg
					Ser
					Leu
					His
					Cys>
660	670	680	690	700	
AAG	CCT	GAG	GAA	GCC	CTT
Lys	Pro	Glu	Glu	Ala	Leu
710	720	730	740	750	760
CAG	GCT	GCC	ACC	TCC	TCC
Gln	Ala	Ala	Thr	Ser	Ser
770	780	790	800	810	
CCC	ACT	GCT	GGG	TCA	ACA
Pro	Thr	Ala	Gly	Ser	Thr
					Asp
					Pro
					Pro
					Gln
					Ser
					Pro
					Gln
					Gly
					Ala
					Ser
					Ala
					Phe>

SUBSTITUTE SHEET (RULE 26)

(Sheet 2 of 3)

```

820      830      840      850      860      870
.      .      .      .      .      .
CCC ACT ACC ATC AAC TTC ACT CGA CAG AGG CAA CCC AGT GAG GGT TCC AGC AGC
Pro Thr Thr Ile Asn Phe Thr Arg Gln Arg Gln Pro Ser Glu Gly Ser Ser Ser>

      880      890      900      910      920
      .      .      .      .      .
CGT GAA GAG GAG GGG CCA AGC ACC TCT TGT ATC CTG GAG TCC TTG TTC CGA GCA
Arg Glu Glu Glu Gly Pro Ser Thr Ser Cys Ile Leu Glu Ser Leu Phe Arg Ala>

      930      940      950      960      970
      .      .      .      .      .
GTA ATC ACT AAG AAG GTG GCT GAT TTG GTT GGT TTT CTG CTC CTC AAA TAT CGA
Val Ile Thr Lys Lys Val Ala Asp Leu Val Gly Phe Leu Leu Leu Lys Tyr Arg>

980      990      1000      1010      1020      1030
.      .      .      .      .      .
GCC AGG GAG CCA GTC ACA AAG GCA GAA ATG CTG GAG AGT GTC ATC AAA AAT TAC
Ala Arg Glu Pro Val Thr Lys Ala Glu Met Leu Glu Ser Val Ile Lys Asn Tyr>

      1040      1050      1060      1070      1080
      .      .      .      .      .
AAG CAC TGT TTT CCT GAG ATC TTC GGC AAA GCC TCT GAG TCC TTG CAG CTG GTC
Lys His Cys Phe Pro Glu Ile Phe Gly Lys Ala Ser Glu Ser Leu Gln Leu Val>

1090      1100      1110      1120      1130      1140
.      .      .      .      .      .
TTT GGC ATT GAC GTG AAG GAA GCA GAC CCC ACC GGC CAC TCC TAT GTC CTT GTC
Phe Gly Ile Asp Val Lys Glu Ala Asp Pro Thr Gly His Ser Tyr Val Leu Val>

      1150      1160      1170      1180      1190
      .      .      .      .      .
ACC TGC CTA GGT CTC TCC TAT GAT GGC CTG CTG GGT GAT AAT CAG ATC ATG CCC
Thr Cys Leu Gly Leu Ser Tyr Asp Gly Leu Leu Gly Asp Asn Gln Ile Met Pro>

      1200      1210      1220      1230      1240
      .      .      .      .      .
AAG ACA GGC TTC CTG ATA ATT GTC CTG GTC ATG ATT GCA ATG GAG GGC GGC CAT
Lys Thr Gly Phe Leu Ile Ile Val Leu Val Met Ile Ala Met Glu Gly Gly His>

1250      1260      1270      1280      1290      1300
.      .      .      .      .      .
GCT CCT GAG GAG GAA ATC TGG GAG GAG CTG AGT GTG ATG GAG GTG TAT GAT GGG
Ala Pro Glu Glu Glu Ile Trp Glu Glu Leu Ser Val Met Glu Val Tyr Asp Gly>

      1310      1320      1330      1340      1350
      .      .      .      .      .
AGG GAG CAC AGT GCC TAT GGG GAG CCC AGG AAG CTG CTC ACC CAA GAT TTG GTG
Arg Glu His Ser Ala Tyr Gly Glu Pro Arg Lys Leu Leu Thr Gln Asp Leu Val>

1360      1370      1380      1390      1400      1410
.      .      .      .      .      .
CAG GAA AAG TAC CTG GAG TAC CGG CAG GTG CCG GAC AGT GAT CCC GCA CGC TAT
Gln Glu Lys Tyr Leu Glu Tyr Arg Gln Val Pro Asp Ser Asp Pro Ala Arg Tyr>

      1420      1430      1440      1450      1460
      .      .      .      .      .
GAG TTC CTG TGG GGT CCA AGG GCC CTC GCT GAA ACC ACC TAT GTG AAA GTC CTT
Glu Phe Leu Trp Gly Pro Arg Ala Leu Ala Glu Thr Ser Tyr Val Lys Val Leu>

      1470      1480      1490      1500      1510
      .      .      .      .      .

```

(Sheet 3 of 3)

GAG TAT GTG ATC AAG GTC AGT GCA AGA GTT CGC TTT TTC TTC CCA TCC CTG CGT  
 Glu Tyr Val Ile Lys Val Ser Ala Arg Val Arg Phe Phe Phe Pro Ser Leu Arg>

1520 1530 1540 1550 1560 1570  
 GAA GCA GCT TTG AGA GAG GAG GAA GAG GGA GTC TGA GCATG AGTTG CAGCC AAGGC  
 Glu Ala Ala Leu Arg Glu Glu Glu Glu Gly Val>

1580 1590 1600 1610 1620 1630  
 CAGTG GGAGG GGGAC TGGGC CAGTG CACCT TCCAG GGCCG CGTCC AGCAG CTTCCT CCTGC

1640 1650 1660 1670 1680 1690  
 CTCGT GTGAC ATGAG GCCCA TTCTT CACTC TGAAG AGAGC GGTC A GTGTT CTCAG TAGTA

1700 1710 1720 1730 1740 1750  
 GGTTC CTGTT CTATT GGGTG ACTTG GAGAT TTATC TTTGT TCTCT TTTGG AATTG TTCAA

1760 1770 1780 1790 1800 1810  
 ATGTT TTTTT TTAAG GGATG GTTGA ATGAA CTTCA GCATC CAAGT TTATG AATGA CAGCA

1820 1830 1840 1850 1860 1870  
 GTCAC ACAGT TCTGT GTATA TAGTT TAAGG GTAAG AGTCT TGTGT TTTAT TCAGA TTGGG

1880 1890 1900 1910 1920 1930  
 AAATC CATTG TATTT TGTGA ATTGG GATAA TAACA GCAGT GGAAT AAGTA CTTAG AAATG

1940 1950 1960 1970 1980 1990  
 TGAAA AATGA GCAGT AAAAT AGATG AGATA AAGAA CTAAA GAAAT TAAGA GATAG TCAAT

2000 2010 2020 2030 2040 2050  
 TCTTG CCTTA TACCT CAGTC TATTC TGTA AATTT TTAAG GATAT ATGCA TACCT GGATT

2060 2070 2080 2090 2100 2110  
 TCCTT GGCTT CTTTG AGAAT GTAAG AGAAA TTAAG TCTGA ATAAA GAATT CTTCCT TGTTT

2120 2130 2140 2150 2160 2170  
 ACTGG CTCTT TTCTT CTCCA TGCAC TGAGC ATCTG CTTTT TGGAA GGCCC TGGGT TAGTA

2180 2190 2200 2210 2220 2230  
 GTGGA GATGC TAAGG TAAGC CAGAC TCATA CCCAC CCATA GGGTC GTAGA GTCTA GGAGC

2240 2250 2260 2270 2280 2290  
 TGCAG TCACG TAATC GAGGT GGCAA GATGT CCTCT AAAGA TGTAG GGAAA AGTGA GAGAG

2300 2310 2320 2330 2340 2350  
 GGGTG AGGGT GTGGG GCTCC GGGTG AGAGT GGTGG AGTGT CAATG CCCTG AGCTG GGGCA

2360 2370 2380 2390 2400 2410  
 TTTTG GGCTT TGGGA AACTG CAGTT CCTTC TGGGG GAGCT GATTG TAATG ATCTT GGGTG

2420  
 GATCC

Figure 2

<u>GB</u>	MSLEQRS <sup>LHCK</sup> PEEAL <sup>EAQ</sup> Q <sup>ALGLVC</sup> VQ <sup>AA</sup> TSS <sup>SSPLVLG</sup> TL <sup>EEV</sup> PTAG <sup>STD</sup> PP <sup>QSPQ</sup> Q <sup>ASAF</sup> TT <sup>INF</sup> TR <sup>QRP</sup> SEG <sup>SSS</sup> REEG <sup>PTSC</sup> ILES <sup>LFRA</sup> VI <sup>TKK</sup> VAD <sup>LVG</sup> FL <sup>LLKY</sup> RARE <sup>PVT</sup> KAE <sup>MLES</sup> VI <sup>KNY</sup> KH <sup>CFPE</sup> IFG <sup>KASE</sup> SL <sup>QLV</sup> FG <sup>IDV</sup> KEAD <sup>P</sup> TGH <sup>SYVL</sup> VT <sup>CLGL</sup> SYD <sup>GLLG</sup> DN <sup>QIMP</sup> KT <sup>GFLI</sup> IV <sup>LVMI</sup> AME <sup>GCHA</sup> PEEE <sup>IWEEL</sup> SV <sup>MEV</sup> YD <sup>GREH</sup> SA <sup>YGE</sup> PR <sup>KLLT</sup> QD <sup>LV</sup>
<u>GB</u>	Q E K Y L E Y G R C R T V I P H A M S S C CAG GAA AAG TAC CTG GAG TAC -GGC AGG TGC CGG ACA GTG ATC CCG CAC GCT ATG AGT TCC TGT CAG GAA AAG TAC CTG GAG TAC CCG CAG GTG CCG GAC GAT CCC GCA CGC TAT GAG TTC CTG
<u>CY</u>	Q E K Y L E Y R Q V P D S D P A R Y E F L
<u>GB</u>	GVQGPS <sup>LKPAM</sup> *
<u>CY</u>	WGPRA <sup>LAETS</sup> YV <sup>VKV</sup> LE <sup>YVI</sup> KV <sup>SA</sup> R <sup>VR</sup> PP <sup>PS</sup> LR <sup>EAA</sup> LR <sup>EEEE</sup> GV





Fig. 3A

## Peptide Screening: MAGE/A3.






Fig. 3C



## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US94/08721

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :Please See Extra Sheet.

US CL :Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 424/277.1; 435/69.3, 240.2, 252.3, 254, 320.1; 530/324, 327, 328, 828; 935/12

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

Please See Extra Sheet.

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SCIENCE, Volume 254, issued 13 December 1991, P. Van Der Bruggen et al., "A Gene Encoding an Antigen Recognized by Cytolytic T Lymphocytes on a Human Melanoma", pages 1643-1647, see entire document.	1-8
X	INTERNATIONAL JOURNAL OF CANCER, Volume 52, issued 1992, F. Brasseur et al., "Human Gene MAGE-1, Which Codes for a Tumor Rejection-Antigen, is Expressed by Some Breast Tumors", pages 839-841, see entire document.	1-8

☐ Further documents are listed in the continuation of Box C.
 ☐ See patent family annex.

Special categories of cited documents:	
*A* document defining the general state of the art which is not considered to be of particular relevance	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
*E* earlier document published on or after the international filing date	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
*L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
*O* document referring to an oral disclosure, use, exhibition or other means	
*P* document published prior to the international filing date but later than the priority date claimed	*G* document member of the same patent family

Date of the actual completion of the international search 05 OCTOBER 1994	Date of mailing of the international search report 25 NOV 1994 -
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231	Authorized officer ANTHONY C. CAPUTA, Ph.D. <i>R. Kuyza for</i>
Facsimile No. (703) 305-3230	Telephone No. (703) 308-0196

# INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US94/08721

## A. CLASSIFICATION OF SUBJECT MATTER:

IPC (6):

A61K 38/00, 39/00; C07H 21/04; C12N 1/19, 1/21, 5/10, 15/00, 15/70, 15/79; C12P 21/06

## A. CLASSIFICATION OF SUBJECT MATTER:

US CL :

424/277.1; 435/69.3, 240.2, 252.3, 254, 320.1; 530/324, 327, 328, 828; 935/12

## B. FIELDS SEARCHED

Electronic data bases consulted (Name of data base and where practicable terms used):

APS, CA, BIOSIS, MEDLINE, EMBASE, WPIDS,

search terms: fikes, j?, livingston, b?, sette, a?, mage 1, mage, terminal, end, carboxyl, C terminal,